

Techniques de réécriture

TD n°3 : Equational theories & completion

Definition. Fix an alphabet \mathcal{F} and a countably infinite set of variables \mathcal{X} . An identity is a pair $(s, t) \in T(\mathcal{F}, \mathcal{X}) \times T(\mathcal{F}, \mathcal{X})$. Fix a set E of identities. We define \approx_E , the induced equality, by the following rules :

$$\frac{(s, t) \in E}{s \approx_E t} \quad \frac{}{t \approx_E t} \quad \frac{s \approx_E t}{t \approx_E s} \quad \frac{s \approx_E t \quad t \approx_E u}{s \approx_E u}$$

$$\frac{s \approx_E t}{s\sigma \approx_E t\sigma} \quad \frac{s_1 \approx_E t_1 \cdots s_n \approx_E t_n}{f(s_1, \dots, s_n) \approx_E f(t_1, \dots, t_n)}$$

Definition. We are interested in the two following problems :

(Word Problem for E) (with E fixed) :

Instance : two terms s and t .

Question : $s \approx_E t$?

(Word Problem) :

Instance : a finite set E of identities and two terms s and t .

Question : $s \approx_E t$?

Exercise 1 :

- 1) Do you know a finite set of identities for which the word problem is undecidable ?
- 2) What can you say about **(Word Problem)** ?

Exercise 2 :

- 1) Given a set of identities E , prove that $R = E \cup E^{-1}$ is confluent.
- 2) Prove that the **(Confluence Problem)** :

Instance : a finite rewrite system R .

Question : Is R confluent ?

is undecidable.

Definition. We say that two sets of identities E and E' are equivalent if $\approx_E = \approx_{E'}$.

Exercise 3 :

What is the behavior of the basic completion procedure on the following set of identities (with the suitable reduction order) :

- $\{(x * (y + z), (x * y) + (x * z)), ((u + v) * w, (u * w) + (v * w))\}$ and $>$ the lpo with $* > +$
- $\{(x + 0, x), (x + s(y), s(x + y))\}$ and $>$ the kbo with $s > +$ and weight 1 for all variables and symbols
- $\{(f(g(f(x))), x)\}$
- $\{(f(g(f(x))), f(g(x)))\}$
- $\{((x * y) * (y * z), y)\}$ with any simplification order

Algorithm 1 Basic completion procedure

Require: A finite set E of identities and a reduction order $>$

Ensure: A finite convergent rewrite system R equivalent to E if the procedure terminates successfully, FAIL if the procedure terminates unsuccessfully

```
1: if there exists  $(s, t) \in E$  such that  $s \neq t$ ,  $s \not\approx t$  and  $t \not\approx s$  then
2:   terminates with output FAIL
3: else
4:    $i := 0$ 
5:    $R_0 := \{(l, r) \mid (l, r) \in E \cup E^{-1} \wedge l > r\}$ 
6: end if
7: repeat
8:    $R_{i+1} := R_i$ 
9:   for all  $(s, t) \in CP(R_i)$  do
10:    Reduce  $s$  and  $t$  to some  $R_i$ -normal forms  $\tilde{s}$  and  $\tilde{t}$ 
11:    if  $\tilde{s} \neq \tilde{t} \wedge \tilde{s} \not\approx \tilde{t} \wedge \tilde{s} \not\approx \tilde{t}$  then
12:      terminates with output FAIL
13:    end if
14:    if  $\tilde{s} > \tilde{t}$  then
15:       $R_{i+1} := R_{i+1} \cup \{(\tilde{s}, \tilde{t})\}$ 
16:    end if
17:    if  $\tilde{t} > \tilde{s}$  then
18:       $R_{i+1} := R_{i+1} \cup \{(\tilde{t}, \tilde{s})\}$ 
19:    end if
20:  end for
21:   $i := i + 1$ 
22: until  $R_i = R_{i+1}$ 
23: return  $R_i$ 
```

Exercise 4 :

You can admit the correction of the procedure.

- 1) Prove that the Huet's completion procedure always successfully terminates on finite sets of ground identities and $>$ a reduction order that is total on ground terms.
- 2) Deduce that **(Word Problem)** is decidable for finite sets of ground identities.

Exercise 5 :

- 1) Apply the Huet's completion procedure on $\{(f(x, y), h(x, c)), (f(x, y), h(c, y)), (h(x, c), c)\}$ and the lpo with $f > h > c$.
- 2) Prove that $R = \{f(x, y) \rightarrow c, h(c, y) \rightarrow c, h(x, c) \rightarrow c\}$ is equivalent to the set of identities of question 1.
- 3) Notice that R terminates. Prove this termination using the lpo of question 1. Deduce that R is convergent.

Exercise 6 :

- 1) Prove that both completion procedures may fail on

$$\{(h(x, y), f(x)), (h(x, y), f(y)), (g(x, y), h(x, y)), (g(x, y), a)\}$$

and the lpo with $g > h > f > a$.

- 2) How can we avoid the failure?

Algorithm 2 Huet's completion procedure

Require: A finite set E of identities and a reduction order $>$

Ensure: A finite convergent rewrite system R equivalent to E if the procedure terminates successfully, FAIL if the procedure terminates unsuccessfully

```
1:  $R_0 := \emptyset$ ;  $E_0 := E$ ;  $i := 0$ 
2: while  $E_i \neq \emptyset$  or there is an unmarked rule in  $R_i$  do
3:   while  $E_i \neq \emptyset$  do
4:     Choose an identity  $(s, t) \in E$ 
5:     Reduce  $s$  and  $t$  to some  $R_i$ -normal forms  $\tilde{s}$  and  $\tilde{t}$ 
6:     if  $\tilde{s} = \tilde{t}$  then
7:        $R_{i+1} := R_i$ ;  $E_{i+1} := E_i \setminus \{(s, t)\}$ ;  $i := i + 1$ 
8:     else
9:       if  $\tilde{s} \not\approx \tilde{t} \wedge \tilde{s} \not\approx \tilde{t}$  then
10:        terminates with output FAIL
11:      else
12:        let  $l$  and  $r$  such that  $\{l, r\} = \{\tilde{s}, \tilde{t}\}$  and  $l > r$ 
13:         $R_{i+1} := \{(g, \tilde{d}) \mid (g, d) \in R_i \wedge g \text{ cannot be reduced with } l \rightarrow r \wedge \tilde{d} \text{ is a } R_i \cup \{(l, r)\}\text{-normal form of } d\} \cup \{(l, r)\}$ 
14:         $(l, r)$  is not marked and  $(g, \tilde{d})$  is marked in  $R_{i+1}$  iff  $(g, d)$  is in  $R_i$ 
15:         $E_{i+1} := (E_i \setminus \{(s, t)\}) \cup \{(g', d) \mid (g, d) \in R_i \wedge g \text{ can be reduced to } g' \text{ with } l \rightarrow r\}$ 
16:         $i := i + 1$ 
17:      end if
18:    end if
19:  end while
20:  if there is an unmarked rule in  $R_i$  then
21:    let  $(l, r)$  be such a rule
22:     $R_{i+1} := R_i$ 
23:     $E_{i+1} := \{(s, t) \mid (s, t) \text{ is a critical pair of } (l, r) \text{ with itself or with a marked rule in } R_i\}$ 
24:     $i := i + 1$ 
25:    Mark  $(l, r)$ 
26:  end if
27: end while
28: return  $R_i$ 
```
